

Volcanic risk to marine infrastructure and shipping

 Paul Cragg^α,  Pete Rowley^{*β}, and  Samuel Mitchell^β

^α Unaffiliated.

^β School of Earth Sciences, University of Bristol, Bristol, UK.

ABSTRACT

Exploration of volcanic risk over recent decades has helped garner a detailed understanding of the vulnerabilities and processes at work in the terrestrial setting, but relatively little is understood about volcanic risk at sea, despite our increasing reliance on surface shipping, energy and resource infrastructure, and submarine telecoms cables. This is rooted in (1) a lack of understanding of the marine volcanic processes themselves, (2) a lack of knowledge regarding how these processes interact with existing use and infrastructure at sea, and (3) no effective quantification of the degree of exposure and therefore vulnerability that different elements of our trade and industry experience to these different hazard processes. Addressing these concerns must be a priority for the volcanological and maritime communities, as well as the users and stakeholders. This work aims to shine a light on some of the processes, interactions and major vulnerabilities which we are currently exposed to as a globally connected society. We focus primarily on exploring the primary hazards and detailing the risk posed to shipping and port infrastructure. Some first order constraints are explored to provide context for the scale of the issues at hand. Finally, we suggest 6 initial goals for disaster risk reduction which require a full spectrum of research, policy, and behavioural developments and improvements.

KEYWORDS: Volcanology; Marine hazard; Risk; Shipping; Infrastructure.

1 INTRODUCTION

The rapid development of subsea infrastructure and the growth in international shipping has vastly increased exposure to little-studied marine volcanic hazards. Quantification of volcanic risk in the terrestrial environment has been an area of considerable research and progress in the last 50 years [Marzocchi and Woo 2009; Brown et al. 2015; Loughlin et al. 2015]. Exposure of infrastructure and populations living on or near active volcanoes to tephra fall, pyroclastic density current, lahars, and lava flow activity is well recognised [e.g., Barsotti et al. 2018; Bevilacqua et al. 2022; Nagai and Nakada 2022; Jenkins et al. 2024]. Experimental and numerical modelling approaches have been used to explore the extent of this exposure to different types and scales of volcanic activity [e.g., Thouret et al. 2023; Aravena et al. 2024; Engwell et al. 2024]. These assessments, however, are usually constrained to sub-aerial volcanoes and rarely extend offshore, despite considerable known impacts already experienced around submarine, ocean island, and near-shore volcanoes, and up to thousands of kilometres from volcanic sources. A good example of hazard being mapped offshore is Waythomas and Waitt [1998], which explores offshore tephra fall, pyroclastic density current, debris avalanche, and directed blast hazards from Augustine volcano. In terms of submarine examples, Verolino et al. [2024] explore seamount hazard in the Indian Ocean, and Maeno [2025] focusses on the Kolumbo field.

There have often been papers resulting from recent or ongoing events which explore individual hazard processes, for example, flank collapse-generated tsunamis [Legros and Druitt 2000; Milia et al. 2003; Le Friant et al. 2009; Cas et al. 2024], propagation and entry of pyroclastic density currents into the sea [Di Roberto et al. 2014; Giordano and De Astis 2020; Andronico et al. 2021], and pumice raft formation [Bryan et al.

2012; Iskandar et al. 2023]. Studies have also resulted from the exploration of historical deposits during expeditions [e.g., Satow et al. 2023; Druitt et al. 2024; Walding et al. 2026]. General hazards posed to marine infrastructure by pumice rafts have been discussed previously [Jutzeler et al. 2014], as well as the identification of some critical choke points where low magnitude volcanic activity could disrupt aerial, maritime, transport and trade, and submarine infrastructure [Mani et al. 2021]. The vulnerability of submarine cable infrastructure has been the focus of recent work, highlighting the 2022 Hunga Tonga-Hunga Ha'apai eruption [Clare et al. 2023], with cable break examples dating back to Krakatau in 1883 [Clare et al. 2025]. Recent work has explored the impact of volcanic ash to both shipping and infrastructure in the context of inland waterways in Patagonian lakes following activity at Cordón Caulle, Chaitén, and Calbuco [Salgado et al. 2023]. This record of impacts establishes a broad range of damage types done to ship systems and port infrastructure in the case of small vessels, ranging from capsizing to oil and fuel contamination, abrasive hull damage, corrosion, and failure of electrical systems.

In this work we take the insights from these existing works, combined with geospatial analyses of marine traffic and port infrastructure, and insight into ship operations and mechanisms to present an overview of maritime volcanic hazards. We briefly explore the exposure of shipping and the magnitude of the industry presently exposed. We then discuss the volcanic hazard processes which may pose the greatest threat to shipping and infrastructure (Figure 1) before looking at specific vulnerabilities that ships have in relation to these processes. Finally, we suggest some priority areas for the natural hazard community to focus on based on our current understanding of frequency, range, and impact of different hazards.

*✉ pete.rowley@bristol.ac.uk




Figure 1: Schematic of the major volcanic hazards with the potential to impact the marine and submarine realm. Ballistic bombs are not illustrated. The impact of lava flows to shoreline activity is also not included.

2 THE GROWTH OF SHIPPING AND ITS INTERSECTION WITH VOLCANISM

Approximately 90 % of global trade is reliant on marine shipping, with the volume of shipping traffic forecast to more than double from its 2018 values by 2050 [Sardain et al. 2019]. The forecasts for the magnitude of shipping growth, which vary depending on the controls on modelled growth rates (ranges between a 63–210 % increase by 2050), cover a broad variety of economic, geopolitical, technical, and environmental forecasts [Walsh et al. 2019; International Maritime Organization 2020]. While some of these projections predate the 3.8 % reduction in shipping in 2020 as a result of the global COVID19 pandemic, subsequent rebound of 2.4 % growth in 2023 is forecast to continue at ~2 % through to 2028 [United Nations Trade and Development 2023].

Many key shipping routes pass within 100 km of active volcanic systems (Figure 2), and there is widespread documented evidence of ships being affected by eruptive phenomena throughout recent history, for example at the eruptions from Myojin-sho in 1792 [Morimoto and Osaka 1970], Krakatau in 1883 [Simkin and Fiske 1983], Montagne Pelée in 1902 [Johnson and Threlfall 1985; Tanguy 1994; Scarth 2002; Zebrowski 2002], and Rabaul Caldera in 1937 [Johnson and Threlfall 1985] and 1994 [Global Volcanism Program 1994]. Impact to ships from marine volcanic hazards date back to at least 1650 BCE during the Late Bronze Age eruption of Santorini [Evans and McCoy 2020; Karátson et al. 2020]. In Simkin and Fiske [1983] there are nearly 100 ships mentioned regarding encounters with floating pumice and tephra fall as a result of the Krakatau eruption in 1883, with notable ash loading on a ship deck reported over 1000 km away. Shipping losses, while less common, have most recently been documented due to the the Hunga Tonga-Hunga Ha'apai eruption in 2022, causing vessel losses as far afield as Chile and Japan due to the far-reaching volcanic-induced tsunamis [Terry et al. 2022; The Guardian 2022; World Bank 2022]. A summary of

reported shipping impacts and losses from specific eruptions is provided in Table 1.

The exposure of maritime trade and transport goes further than simply shipping routes, however, as a substantial number of ports also lie within range of many of the hazards identified in Table 2. Figure 2B-E highlight the co-location of major ports in some of the worlds high traffic exposure. A spatial analysis exploring how many Holocene volcanoes lie within 100 km of each of the 856 ports within the World Port Index database reveals that many have at least some exposure to local volcanism (Figure 3). Over 100 million Twenty-Foot Equivalent Units (TEU) of cargo capacity are exported annually from ports with 3 or more volcanoes within 100 km, rising to over 300 million TEU when considering those close to a single Holocene volcano. These numbers only consider containerised transport. Acajutla and San Salvador ports in El Salvador both lie within 100 km of 19 different volcanoes, and 6 of the top 50 cargo ports lie within range of at least one: Yokohama (Japan ~27 M TEU), Piraeus (Greece ~25 M TEU), Barcelona (Spain ~23 M TEU), Tokyo (Japan 21 M TEU), Gioia Tauro (Italy ~17 M TEU), and Manzanillo (Mexico ~15 M TEU). Ports with smaller outflows still exceeding 10 M TEUs include Balboa (Panama), Jakarta (Indonesia), Vancouver (Canada), and Taipei (Taiwan). Note that the available dataset only includes dry cargo and does not include import volumes, which would undoubtedly modify the sequencing. We do not capture the overall picture of trade exposure here but hope to demonstrate the order of magnitude of the issue.

Of the regional data mapped in Figure 2, it is worth highlighting that over 36 % of the 152 identified ports lie in just three countries. Thirty-two of Japan's 35 listed ports lie within 100 km of at least one volcano. Thirteen of 15 ports in the Philippines, and 10 of the 29 ports in Indonesia are similarly exposed.

In addition to the direct threat to vessels, crews, and passengers, the impacts of submarine or near-shore volcanic activity (Figure 1) may include navigation blockage and disruption.




Figure 2: [A] Global map of active volcanoes (white triangles), shipping intensity (orange-red), and observed pumice rafts (arrows and dates) over the last 200 years highlighting substantial overlaps in key areas including [B] the Mediterranean, [C] the Sea of Japan, [D] the Molucca Sea, and [E] the Java, Flores, and Savu Seas. The shipping intensity data [World Bank Group 2021] is derived from hourly Automatic Identification System (AIS) positions received between January 2015 and February 2021. The AIS positions may have been transmitted by both moving and stationary ships within each grid cell, therefore it is analogous to the general intensity of shipping activity and not necessarily the volume of traffic. Black circles indicate major ports [National Geospatial-Intelligence Agency 2022; World Bank Group 2023], with relative scale related to volume of cargo handled. White triangles indicate Holocene volcanoes [Global Volcanism Program 2025].

Table 1: Shipping impacts and losses identified in the literature.

Volcano	Year	Shipping affected	Range of reported impact	References
Krakatau	1883	Over 100 vessels	Ash loading reported over 1000 km away, tsunami impacts on coasts over 1000 km away, navigation hazards from floating pumice and ash over wide areas for weeks or months afterward, stranding of large steam vessel 2+ km inland with the loss of all hands, hundreds of ships impacted	Simkin and Fiske [1983]
Rabaul	1937	Pumice loading (including ingress into holds), abrasion by PDC, harbour impassable	Only nearshore reported	Johnson and Threlfall [1985]
Pelée	1902	Estimated 394 total losses by capsizing or fire, including 386 French ships and boats, eight foreign vessels, 647 crew members, and 34 passengers	Nearshore, Pouyer-Quertier, over 12 km from the shore was damaged	Tanguy [1994], Caron [2013], and Musée Frank A Perret [2019-]
Myojin-sho	1952	Loss of observation ship and all crew (29)	Proximal ('a distance convenient for observations')	Minakami [1956]
Chaitén	2008	Damage and abrasion to deck surfaces, interruption to air supplies, impacts on engine cooling, performance of water jets, damage and disruption to port access and machinery	Up to 100 km	Salgado et al. [2023]
Cordón Caulle	2011–2012	Capsizings, pitch and stability issues, damage and abrasion to deck surfaces, contamination of fuel and oil, effect on electrical systems, interruption to air supplies, impacts on engine cooling, performance of water jets, damage and disruption to port access and machinery	Severe impacts up to 100 km	Salgado et al. [2023]
Calbuco	2015	Damage and abrasion to deck surfaces, interruption to air supplies, impacts on engine cooling, performance of water jets, ash ingress into ship interiors, damage and disruption to port access and machinery	Up to 200 km	Salgado et al. [2023]
Hunga Tonga-Hunga Ha'apai	2022	28 boats capsized, sunk or floated away in Southern Japan, capsized boat in Santa Barbara, USA, some boats sunk in New Zealand, many boats and ports damaged around the Pacific	Over 8000 km	World Bank [2022]
Hekla	1845	Ash landing on ships	820 km - ash loading on ship of 0.03 kg m ⁻²	Thorarinnsson [1950]

tion, resulting in economic impacts at anything from local to global scales, depending on the degree and location of disruption. Exploration of the impacts of non-volcanic navigational blockages, and modelling of global shipping movements has revealed that restriction at key choke points can substantially reroute global shipping [Pratson 2023] and that container and petrochemical tankers are particularly impacted [Lee and Wong 2021].

The potential delays caused by navigational issues due to volcanism are of particular concern given the high vulnerability of various parts of the shipping network to cascading failure as a result of port throughput and the link strength between various nodes on that network [Liupeng et al. 2024; Xu et al. 2024]. Arriving at ports out of sequence can result in further delays and knock-on effects impacting not only the ongoing activity of the delayed ship, but all associated trade goods and supply networks, as well as the scheduling of other vessels. The closure of ports provides an even higher risk to the trade network, whereby re-routing of traffic to alternative destinations destabilises the system faster than simple delays [Xu et al. 2022].

It is important to note that the shipping industry is not solely comprised of commercial trade of goods, fuel, etc., but also comprises tourism and leisure (cruise ships), research expeditions, private vessels, merchant and military vessels, and the global fishing/aquaculture industry. The distribution and density of these individual industries may vary due to location, distance to shore, trade routes, access to territorial waters, territorial disputes, culture, geopolitics, climate, local ecosystems, seasonal changes, or national affluence. This variety of factors make the assessment of volcanic risk to shipping a globally complex issue.

3 HAZARD PROCESSES

Recorded shipping losses have been attributed to a range of processes, including capsizing due to volcanically-induced tsunamis [Imamura et al. 2022], inundation by pyroclastic density currents [Guibert et al. 2018], and damage by tephra fall [Salgado et al. 2022; Salgado et al. 2023]. Submarine volcanism also produces widespread and poorly monitored pumice rafts (Figure 1), which may be a hinderance to transit, or pose mechanical risks. The risks posed and range of impact of these different hazards varies considerably (Table 2).

Pumice rafts pose perhaps the longest duration and widest dispersed threat. Our estimate of frequency of pumice raft-forming events is very poor, as often these are only identified for scientific recording when the rafts encounter shorelines, possibly months after eruption at remote submarine volcanoes [Bryan et al. 2004; Jutzeler et al. 2014; Redick 2023]. The lifetime of pumice rafts is strongly controlled by the microstructure of the pumice [Mitchell et al. 2021; Nagao et al. 2025], while distribution is governed by the interaction of wind, wave, and ocean currents [Bryan et al. 2004; Jutzeler et al. 2014]. Satellite observation can then be used to back-calculate the tracks and ultimate sources of these rafts. Increasing efforts are being made to use and develop satellite-based detection and tracking to mitigate the impacts of raft encounters




Figure 3: Evaluation of number of volcanoes within 100 km for all ports included in the World Port Index by region. 152 of the 856 listed ports (~18 %) lie within 100 km of at least one Holocene volcano. 67 ports lie within 100 km of 3 or more volcanoes, while two ports in El Salvador are within 100 km of 19 volcanoes. The largest number of exposed ports lie in Japan.

[e.g., using floating boom lines; Whiteside et al. 2021; Zheng et al. 2022; Iskandar et al. 2023; He et al. 2025].

Raft encounters have not only impacted ships [Gass et al. 1963; Nishikawa et al. 2023] but also coastal access and ports [Asano and Nagayama 2021]. For example, the eruption of Fukutoku-Oka-no-Ba submarine volcano caused disruption to coastlines stretching from the Nansei islands to mainland Japan between October–November 2021; two months after the August eruption over 1000 km away [Yoshida et al. 2022; Chang et al. 2023]. The impact of the Chaitén (2008), Cordon-Caulle (2010–2011), and Calbuco (2015) eruptions on Patagonian lakes has provided a rich dataset for the impact of tephra on a shallow water access, as well as impacts on ship systems, including propulsion and structural damage due to abrasion [Salgado et al. 2023].

Tephra fall from eruption columns can be spread over very wide areas, although the distances over which substantial (> 1 mm-thick) deposits can be generated is much more restricted (in the order of 10s to 100s of km from the vent). This still places many high intensity shipping routes within exposed areas as identified by global ash dispersion potential [Jenkins et al. 2015] (Figure 4). Plumes from explosive volcanic eruptions can achieve heights from hundreds of meters to tens of kilometres and disperse ash from local to global scales [Bonadonna and Costa 2013]. Following a number of aviation incidents with ash plume encounters, work has been done to under-




Figure 4: Shipping (Automatic Identification System) AIS data, overlaid with recurrence interval of ashfall exceeding 1 mm thickness [Jenkins et al. 2015] for Southeast Asia, indicating an extensive overlap between high-traffic shipping routes and the arc volcanism ash dispersal hazard for Southeast Asia and the East Pacific. Although this represents the most intense set of choke points and overlap, similar relationships can be observed around the planet, including the Caribbean, Mediterranean, and Red Sea. Holocene eruption data from the Global Volcanism Program [2025].

stand plume transport and the effects of volcanic ash on aircraft [Prata and Tupper 2009; Alexander 2013]. No such work has been completed exploring the impacts of ash ingestion or loading on ship structures and engines.

Pyroclastic density currents are a shorter-range hazard, with typical propagation distances of up to a few tens of kilometres [Freundt 2003]. However, on encountering water they pose a multihazard scenario where several different hazardous processes can occur [e.g., López-Saavedra and Martí 2023]; the more dilute portions of these currents are able to propagate across the water surface and therefore are able to directly impact surface shipping and infrastructure some considerable distance from the shoreline. This was the cause of vessel losses at St Pierre, Martinique in 1902 [Jaggard 1949]. The denser portions of the currents are able to enter the wa-

ter as the flow encounters the shoreline to form subaqueous eruption-fed density currents [Frey et al. 2026]. This has two separate products; the displacement can trigger tsunamis [Bonaccorso et al. 2003; Bouguin et al. 2020; Lipiejkó et al. 2022; Lynett et al. 2022], and the sediment mixes with water to generate submarine volcanoclastic density currents which can propagate for hundreds of kilometres offshore [Trofimovs et al. 2012; Jutzeler et al. 2017]. These currents are not necessarily restricted to subaerial feeding; submarine eruptions which generate steam-charged pumices may also be a source. On encountering cold sea water, the steam within the vesicles rapidly condenses, creating negative pore pressures which then draw in sea water, rapidly densifying the clasts and causing them to sink [Allen et al. 2008]. This process has the potential to feed density currents immediately surround-

ing submarine vents. Such currents pose a substantial risk to subsea infrastructure and were responsible for the breaking of telecommunications cables, most recently during the Hunga Tonga-Hunga Ha'apai eruption in 2022 [Clare et al. 2023; Seabrook et al. 2023]. However, the historical record of these processes causing cable breaks goes back as far as Krakatoa, 1883 [Simkin and Fiske 1983], and Mt Pelée in 1902 [Chrétien and Brousse 1989]. There are at least 11 cases in the historical record where volcanic activity has resulted in submarine cable breaks [Clare et al. 2025].

Volcanogenic tsunamis are also generated in the wake of mass wasting/flank collapse events [Bonaccorso et al. 2003; Giachetti et al. 2012] which, in the case of ocean island volcanoes, can lead to total edifice failure and explosive magma-water interactions as was observed at Krakatau in 1883 [Simkin and Fiske 1983]. These can generate tsunamis greater in scale than those generated by ocean-entering PDCs, as these types of mass wasting generally introduce a larger volume of material to the water very rapidly. The frequency of these events is much lower than eruptive processes; a volcano may take many thousands of individual eruptions to build up, before a collapse removes a substantial proportion of its total mass. The frequency of flank collapse eruptions in subaerial volcanoes corroborates this, with the notable examples of Bezymianny, Russia [Shevchenko et al. 2020], Mount St. Helens, USA [Sarna-Wojcicki et al. 1980], and Anak Krakatau, Indonesia [Williams et al. 2019] in the last century, in line with the estimated average of approximately 4 per century [McGuire 2003]. There can be more directionality in volcanogenic tsunamis compared to tectonically-forced tsunamis. Their generation by displacement by oriented processes like flank collapse [Giachetti et al. 2012; Williams et al. 2019] and entry of volcanoclastic currents [e.g., Ripepe and Lacanna 2024] results in spatially variable wave heights and short wavelengths, similar to those generated by flank collapse and debris-avalanche events [e.g., Williams et al. 2019]. Longer wavelength tsunamis can be caused by atmospheric pressure waves generated by eruptions; these typically have wavelengths and amplitudes which relate directly to water depth, in a way similar to that observed in tectonically-driven tsunami events [Ward and Day 2003; Paris 2015; Grilli et al. 2019]. The eruption of Hunga Tonga-Hunga Ha'apai in 2022 demonstrated multiple tsunamigenic processes, which included this meteogenic process, as well as waves driven by the explosive shock wave and the subsequent collapse of the cavity [Lynett et al. 2022].

Other, less explored hazards are more difficult to speak to with any certainty due to lack of event documentation or real-time detection. Ballistic bombs, for example, are generally only a risk within a few kilometres of a vent, but even so we have recorded encounters such as the tourist boat hit by a decimetre-scale rock ejected during the Kilauea eruption in 2018, generated by a coastal lava-water explosion [BBC News 2018]. Even less documented are the possible risks associated with volcanic gases in the water column or above the water, either from passive degassing, or from eruption driven gases and boiling processes. In such cases there is hypothetically a buoyancy risk due to the decrease in fluid density [Denardo et al. 2001; Hueschen 2010], and the sinking of the Island Queen

over Kick 'em Jenny in 1944 has been attributed to this process [University of the West Indies Seismic Research Centre n.d.(b)]. To our knowledge there are no confirmed reports of such an event involving shipping, and, given the very limited distance over which such a risk would exist, we include it here only for completeness. However, there are recorded instances of the advection volcanic gases from submarine/emergent eruptions proving fatal to communities [e.g., the 1650 AD eruption of Kolumbo, Santorini, that resulted in dozens of fatalities and livestock loss up to 10 km from the eruption source; Fouqué 1879; Nomikou et al. 2012], a potentially significant hazard for ship personnel if exposed with limited risk mitigation.

4 TEPHRA DISPERSAL VULNERABILITIES

4.1 Loss of propulsion

Nearly all marine engines rely upon seawater cooling systems to regulate their running temperatures. Older vessels are able to operate with 50 % reductions in cooling water volume, but newer vessels have lower design overheads and experience issues at as little as 30 % loss of volume of cooling water (Personal Communication, Rolls Royce Marine engineers, Ulsan Shipyard, South Korea. May 2003).

In maritime combustion engines the typical main sea inlet grating for the cooling system restricts the diameter of particles drawn into the sea chest to about 15 mm across the minor axis but could be much greater than this along the major axis. Sand to mud grade particles would initially pass through the system but may eventually start to coat the pipe work and cooling plates [Johnson and Threlfall 1985]. In this scenario, volcanic ash and lapilli will quickly block the internal strainers. Depending on the piece of machinery in question, the coarse filter size in these strainers may be as little as 3 mm mesh. Separate intakes for the auxiliary machinery (generator motors, refrigeration plant, air conditioning, etc.), where fitted, will have finer filters and are likely to quickly become blocked. Keeping these clear will be an added burden to the ship's crew. It is likely that at least some machinery will have to be sacrificed so the main engine and generator motor cooling systems can be maintained.

Diesel engines require an airflow sufficient to ensure a flammable mixture within the combustion chamber. This air comes from the engine room atmosphere which, in turn, is provided by high-capacity ventilation fans. A fine (< 1 mm) mesh filter is generally installed at the air intake to remove solid particles before the air is exhausted into the engine room. If exposed to volcanic ejecta, this filter would soon become clogged with entrained tephra and would need to be removed and cleaned at regular intervals. Increasing tephra fall therefore poses increasing hazards to the crew on deck carrying this out. Marine diesel engines generally operate at temperatures below that required to melt primary volcanic material (> 400 °C), although localised hotspots may exist within cylinders, and some volcanic glasses may already become ductile and semi-molten at temperatures exceeding 400 °C [Giordano et al. 2005]. Some naval vessels are powered by gas turbines, and these can suffer similar problems as experienced by aircraft [Shifler and Choi 2018], where ash is ingested, melted in

Table 2: A catalogue of identified volcanic hazards impacting shipping, summarising approximate range that the hazard may impact and the potential risks to shipping thought to be associated with each. Note this does not include risks to submarine or coastally exposed vehicles, infrastructure, or populations.

Type of volcanic hazard	Proximity to source	Risks to shipping
Lava-water coastal explosions	~1 km	Ship destruction/severe damage, potential loss of life/critical injuries, large ballistics in hull side
Shallow sub-surface explosions	~1 km	Ship destruction/severe damage, potential loss of life/critical injuries, ship instability from wave action
Volcanic gases	Up to 10s of km	Crew inhalation and respiration, possible metal-work corrosion from acidic vapours, bubbles reducing effective buoyancy
Pyroclastic density currents	Up to 10s km	Ship destruction/severe damage, potential loss of life/critical injuries, large ballistics in hull side
Terrestrial and submarine flank collapses	Up to 10s km	Ship destruction/severe damage, potential loss of life/critical injuries, large ballistics in hull side
Tephra (ash/pumice) fallout	Up to 100s km	Damage to electronic equipment, inhalation/respiration of fine ash, instability from wet tephra accumulation, navigation/visibility loss, radar attenuation
Volcanic plumes in atmosphere	Up to 100s km	Disruption to navigation and position fixing, reduced visibility, radar attenuation
Oceanic pumice rafts	Up to 1000s km	Loss of propulsion in engines, need to redirect shipping traffic, mechanical and hull damage, increased fuel costs
Tsunamis and shockwaves	Up to 1000s km	Wave damage, flooding, instability or capsizing, potential loss of life, pressure waves dangerous to hearing and health close to source

the high temperature environment where it can then abrade or stick to components, clogging ventilation, reducing efficiency, or causing outright engine failure [Dunn et al. 1996; Davison and Rutke 2014; Clarkson et al. 2016; Song et al. 2016; Giehl et al. 2017].

4.2 Hazards to stability

The wide dispersal of tephra and its ability to accumulate on ships is a significant potential hazard. For a vessel to float upright, the centre of buoyancy (C_B), centre of gravity (C_G), keel (K), and the metacentre (C_M) must be in a vertical line on the centreline of the vessel, with $KC_M > KC_G > KC_B$ (Figure 5). The metacentre represents a point vertically above the centre of buoyancy, in line with the central axis of the ship. In this case, she is said to have a positive metacentric height (+ $C_G C_M$).

When a vessel is built, the position of her initial metacentre is determined by an inclining experiment and the height of the metacentre above the keel (KC_M) is tabulated for different draughts. To ensure all vessels are stable enough to meet all expected conditions during a voyage a minimum set of criteria are laid down by legislation [e.g., The Merchant Shipping (Load Line) Regulations; UK National Archives 1998], including the minimum $C_G C_M$ that a vessel must have at all stages of her voyage.

Where masses of unknown quantity (both total mass and location) are added to the vessel during transit they must be

accounted for when determining that the ‘departure/arrival’ and ‘at sea’ conditions for the ship meet the minimum legal requirements. These conditions typically occur with excess water and ice loading during a voyage. Excess water mass is generally limited by the catchment area on the surface of the ship and is accounted for both in the minimum criteria mentioned above and particular features within the ship’s construction to allow escape by run off or pumping. The situation that may arise from tephra accumulation is very different, as cohesion and material strength properties come into consideration [Walding et al. 2023; Walding et al. 2025] and instead has more similarities with “icing” in cold climates.

Most vessels experience icing from meeting extreme temperature conditions in moderate to heavy weather. Water spray from the sea hits the cold steelwork of the vessel and deck cargo, where it freezes, and ice accretion occurs. As well as accumulating on the deck and other horizontal surfaces, ice may also build up asymmetrically on vertical surfaces causing the vessel to list as well as loose stability. The International Maritime Organization lays down the minimum allowances required [International Maritime Organization 2016]:

- “All exposed horizontal surfaces are assumed to carry $< 30 \text{ kg m}^2$ of ice.
- Vertical surfaces are assumed to be the lateral area of both sides of the ship above the waterplane and carry less than 7.5 kg m^2 of ice.




Figure 5: Generalised ship stability conditions for [A] stable upright, [B] self-righting, [C] over-loaded upright, and [D] capsizing conditions, considering centres of buoyancy (C_B) and gravity (C_G), and metacentre (C_M) over the keel (K). Buoyancy force (F_B) and weight (F_W) vectors are indicated.

- Other surfaces not included in the above are accounted for by increasing the total of 1 and 2 by 5 % and the calculated moments (mass $\times$ Distance from Keel or Centreline) of these weights by 10 %.

- In this condition the vessel must never become over-loaded."

The magnitude of loading by tephra falls within a broad window of values; while accumulation in proximal areas during large eruptions may reach meters of thickness in a few hours [within a few kilometres of source; e.g., Pyle 1989]; this thins to centimetres then millimetres distally or during smaller eruptions. Dry tephra density generally ranges between 800–2500 kg m⁻³, driven by solid densities in the region of 2200–2700 kg m⁻³ and vesiculation causing open pore space of 0–80 % volume. This pore space is often well connected to the particle surfaces, so tephra will gain water mass in the presence of sea spray or rain, which can lead to even low bulk density tephra achieving bulk densities of over ~2000 kg m⁻³—twice that of ice [Spence et al. 2005; Williams et al. 2021]. At these densities accumulation of just 15 mm of tephra will exceed the ice loading limit of 30 kg m⁻². To put this in context, the 1980 eruption of Mount St. Helens (VEI 5) deposited 15 mm of ash up to 450 km from the vent [Sarna-Wojcicki et al. 1980]. With the tendency of tephra to easily drift in prevailing winds it is likely that any loading on a moving ship will be asymmetrical, a condition which ice loading limit calculations are not suited to, and will likely result in an underestimate of stability risk in these conditions (Figure 5). As wetting occurs, the mechanical properties of the tephra change, so it be-

comes more cohesive and capable of sustaining larger steeper piles [Walding et al. 2023; Walding et al. 2025], enabling even greater loading capacity. Further research is required on the realities of tephra accumulation on ships and how the dynamic load changes under different eruption and weather conditions. Furthermore, we suggest that maritime safety standards need to be updated to ensure that crews who experience tephra loading are aware of the risks associated with this hazard.

4.3 Position fixing and navigation

Cohesive wet volcanic ash provides a clear risk to basic visibility due to its ability to cake windows, as well as the simple visibility reductions which occur from having ash in the air. However, a further problem exists. Marine navigation radar typically operates in either 'S' (10 cm) or 'X' (3 cm) bands. In recent years, radars in these bands have been used extensively to track tephra clouds from volcanic eruptions due to their opacity to radar emissions [Harris et al. 1981; Lacasse et al. 2004; Marzano et al. 2013; Bryan et al. 2017]. The sensitivity of X band radar to tephra (and particularly wet tephra) results in two separate issues; the first is that signal attenuation close to the ship prevents detection of obstacles at range, while the second is that the high reflectance of tephra in the atmosphere leads to signal cluttering. In other words, the effective radar range is reduced, and the signal to noise ratio is vastly impacted. These two effects are demonstrated in Figure 6 where images of an [A] unattenuated, [B] attenuated, and [C] cluttered signal from a ship's navigation radar are shown. We have found no clear evidence that this effect has been recorded or reported, and it is presented here as an item of note. Without further work it is unclear what the scale of this issue might be, the extent or quantity of tephra of what particle sizes or distributions might be most impactful, or how this process might be mitigated or managed if it were to pose navigational issues.

4.4 Electrical systems

It is anticipated that external electrical systems on a ship should not be affected by short term exposure to volcanic products. Electrical equipment and distribution systems on the open deck have at least a watertight and, in many cases, gastight status. Internally, the same standard of protection is not adhered to, and volcanic gases and tephra will be able to gain access to some extent.

The combination of moisture and tephra has been shown to cause serious problems to electrical distribution systems on land [Bebbington et al. 2008; Wardman et al. 2012] and air conditioning units [Gordon et al. 2005]. Radio aerials, especially those used for high frequency transmissions, can be similarly affected by shorting across insulators. In the modern marine environment, there is extensive use of computers, printed circuit boards, and notebooks for critical navigation and control systems. All of these may be affected by exposure to tephra or volcanic gases [Wilson et al. 2012b]. Additionally, a lot of the control equipment is sensitive to temperature and can quickly overheat, making it vulnerable to ash contamination which blocks ventilation [Wilson et al. 2012a]. These items are often kept in specific air-conditioned rooms. Air conditioning




Figure 6: Photographs of ships navigation radar under different conditions. [A] demonstrates strong clear signal from nearby oil rigs, [B] is from the same setting and location, with the radar wrapped in a thin film containing volcanic ash with 30 % moisture to generate attenuation, [C] is unattenuated, at a different time and location, demonstrating extensive noise as a result of a dust storm that has similar reflective properties to volcanic ash.

units have been shown to be vulnerable to failure through tephra coating or blocking of various components. [Salgado et al. \[2022\]](#) note that where internal electrical systems were not sealed, computerised navigation and propulsion control systems experienced flashovers resulting in intermittent issues followed by complete system failure, resulting in an inoperable craft. This was compounded by engine blockage from rafting pumice.

The acidic nature of tephra and volcanic gases may pose an additional threat to systems sensitive to acidic corrosion, although the existing evidence is unclear. Work exploring the impact of tephra leachates on metal roofing systems demonstrated that short term exposure (< 30 days) did not measurably increase corrosion, although these same roofing materials were also resilient to direct application of hydrochloric sulphuric and hydrofluoric acids [\[Oze et al. 2014\]](#). In contrast, impact of acid leachates from tephra fall on water storage systems has been reported in real-world cases from Montserrat [\[Sword-Daniels et al. 2014\]](#). Loss of navigational equipment inside the vessel is unlikely to occur before other problems with navigation, mentioned earlier, become apparent. Loss of control equipment may be a greater problem potentially leading to collision (striking a moving object), allision (striking a stationary object), grounding, or capsizing.

4.5 Health hazards to crew and passengers

Respiration of volcanic ash has been shown, in general, to have little long-term effect on the health of a fit adult [\[Martin et al. 1986; Horwell and Baxter 2006; Stewart et al. 2022\]](#). This is dependent upon the length of exposure and quantity of silica present in ash particles < 4 μm . Irritation and bronchial tract problems are the most likely scenario following short term exposure to tephra particles < 100 μm in size, but inhalation and absorption of particles < 10, 4, and 1 μm can cause different respiratory problems, respectively [\[Stewart et al. 2022\]](#). PM-2.5 face masks can significantly reduce the risks from inhalation of volcanic ash in the eventuality of prolonged exposure. Filtration blocking and/or abrasion within fan motors is

also likely to interfere with air conditioning and refrigeration units on ship, which brings with it potential health and safety impacts for crew and passengers, as well as issues for other ship systems, and, in particular, temperature-sensitive cargo transport. For a vessel caught in an area of tephra fallout, any exposure to passengers or crew is likely to be short term, but ships do not routinely carry appropriate personal protective equipment (PPE), and these risks are not recognised in safety standards set out by the International Maritime Organization (IMO) and the International Labour Organization (ILO).

Additionally, prolonged downwind exposure to volcanic gas (carbon dioxide, sulfur dioxide, and halogen species) may put ship personnel at significant health risk [\[Hansell and Oppenheimer 2004; Horwell and Baxter 2006\]](#). As aforementioned, most deaths associated with the marine-emergent eruption of Kolumbo volcano in 1650 AD, Greece, are attributed to exposure to noxious acidic gases blown over 7 km from the volcanic centre [\[Fouqué 1879\]](#). The interactions of lava and cold seawater at the sea surface can create volcanic “laze” (lava-haze), where chlorine ions dissociate from the salt in seawater and attach to hydrogen ions to form dangerous hydrochloric gases and vapours [\[Resing and Sansone 1999\]](#). However, the risks posed by volcanic gases may also be reduced through appropriate PPE usage.

4.6 Routing and delays

Two non-volcanic cases demonstrate the scale of economic impact when critical shipping lanes and routes are shut down or restricted; the M/V Ever Given blockage of the Suez Canal in 2021 [\[Lee and Wong 2021\]](#) and the blocking of the Strait of Hormuz as a result of military action in 2026. The Suez blockage event lasted just 6 days, yet disrupted supply chains globally, delaying shipments, imposing substantial route diversions and cost the global economy 6–10 billion USD [\[Li 2023\]](#). The impact of the Hormuz blockade is ongoing at the time of writing, so still to be fully evaluated, but approximately 21 % of the world’s oil supply passed through the strait in 2022 [\[Ramadhani and Marzaman 2024\]](#), and the first three weeks of

the blockade reduced the global oil supply by approximately 440 million barrels [Reuters 2026].

Pumice rafts often have lifetimes measured in months, even years [Bryan et al. 2012; Jutzeler et al. 2020], and there are numerous choke points to major shipping lanes which not only pass within short distances of active volcanic centres (Figure 2), but that have also experienced pumice raft activity [e.g., the Sunda Strait, Red Sea, Torres Strait, and others—see Figure 2; Bryan et al. 2012; Pratson 2023]. As documented previously, port closures have resulted from pumice rafts blocking access after numerous recent eruptions, such as at Fukutoku-Oka-Noba in 2021 [Yoshida et al. 2022; Salgado et al. 2023]. The cost of just clearing up the pumice has been estimated at 1.6 B JPY (~8 M GBP in 2026 exchange rates) [Takeuchi et al. 2024]. Such an event in a confined strait with more concentrated shipping would be costly not only in clear up but also in terms of obstructed trade and in repairs/maintenance of impacted vessels.

4.7 Tsunami vulnerabilities

The widespread impacts of tsunamis on not only shipping and ports, but any nearshore infrastructure, has been explored extensively following the 2004 Indian Ocean, 2010 Mentawai, Sumatra, and 2011 Tōhoku earthquakes. Substantial advances in both engineering and early warning systems have been made [Kubo et al. 2005; Bernard and Titov 2015; Borrero et al. 2015], including the deployment of pressure sensor networks, Deep-ocean Assessment and Reporting of Tsunamis (DART) buoys, alongside more co-ordinated sharing of tide gauge and seismic monitoring [Bernard and Titov 2015].

Loss probabilities for different tsunami conditions for ports and moored shipping indicate that even relatively small events can cause substantial damage [Muhari et al. 2015]. While the early warning systems in place for tectonically-forced tsunamis would be effective in identifying volcanologically forced waves, the distribution of sensors is skewed toward subduction systems and is not well placed to provide warning to communities proximal to the volcano. This is exacerbated by the lack of monitoring on most volcanoes.

5 THE WIDER CONTEXT

We are not currently in a position as a community to provide any quantitative measures on frequency or magnitude of events, distributions, or the forces and damage these processes can generate which might require engineering against. Furthermore, a large proportion of the shipping industry still relies on very limited shoreside communication and internet access, with bandwidth in some cases still being prohibited to some ship personnel. A lack of access to rapid information and updates in the event of sudden volcanic activity, which may not be translated through VHF-SRC radio channels, may escalate risk and inhibit appropriate response. Increasing access to satellite internet constellations is already helping to reduce communications issues. Without immediate communication, seafarers are reliant on the guidance documentation that is provided to ships captains. The Mariners Handbook [UK Hydrographic Office 2009] provides general information for seafarers on dealing with natural hazards they may face. It

devotes less than a quarter of a page to volcanic eruptions and only deals with submarine eruptions; about the same amount of space is used to discuss earthquakes, while four pages are devoted to ‘Tropical Revolving Storms’. This section is not mentioned in the Index under ‘Volcano’ but is indexed under ‘Izu Shoto’ (the volcanic chain of islands running southeast from Honshu, Japan) or sub-indexed under ‘Seabed’. Given the severity of possible outcomes outlined here and the rapidity of volcanic hazard development, this bypassing of clarity and signage in the immediately available documentation on ship should raise some concern.

6 SUGGESTED RISK REDUCTION ACTIONS

Improving training and onboard reference materials – IMO and ILO manuals and training should be updated with guidance regarding how to respond to nearby volcanic activity, including training on appropriate PPE, ship management protocols, and awareness of hazards to the ship, personnel, and navigation.

Establishing of temporary exclusion and risk zones during eruption watches and warnings – Implementing predetermined local or regional restrictive zones to maritime traffic, as seen with the active submarine volcanoes Kick ‘em Jenny in the Lesser Antilles [University of the West Indies Seismic Research Centre n.d.(a)] and Mentawai (Kingdom of Tonga). Limits on zone boundaries and implementation should be advised by local observatory scientists and probabilistic modelling of eruption scenarios.

Formalising new networks and communications to notify ships of potential hazards – There is currently only a limited system in place to routinely notify and advise maritime traffic to all nearby volcanic hazards. A system similar to the Volcanic Ash Advisory Centres (VAACs) for aviation traffic systems could allow pre-emptive and more appropriate actions in response to developing volcanic crises.

Expansion of volcanic hazard and risk maps to offshore – Some hazards can be mapped offshore due to little change in the physics and processes controlling their distribution (e.g., ashfall and advection of volcanic gases). Other high-risk hazards, such as pyroclastic density currents (PDCs) and volcanic flank collapse, both subaerial and submarine, have poorly constrained interactions with water and this, alongside the resulting submarine volcanoclastic density currents, require detailed experimentation and numerical modelling work to better constrain their behaviours [Furst et al. 2023]. Modelling of potential tsunami wavefront heights, particularly closer to shoreline around near-shore ramps, is also imperative to ship safety and stability.

Identifying shipping lanes of increased risk and vulnerability – There are currently no studies that quantitatively assess the exposure and vulnerability of specific shipping lanes or areas of high marine traffic through industry, fishing, aquaculture, and/or tourism. This requires research that combines both global AIS datasets with known frequency, duration, and intensity of eruptions from proximal volcanic systems. This is certainly an area worthy of attention where the data is available.

Improved knowledge of vulnerable ship systems to different hazards – In this work we have outlined some ship systems likely to be impacted by different volcanic hazards, but the extent of these systems' vulnerability is unknown. We have limited published work which looks directly at ship systems impacted by volcanic activity [e.g., Salgado et al. 2023]. Further work is needed to explore both the processes which impact the systems and the thresholds for safe operation.

Improved knowledge of hazardous coastal/marine volcanic centres – Recent, high-impact eruptions have encouraged a surge of interest in shallow marine and marine-emergent volcanism. However, there are still many large knowledge gaps in this area of research, most critically: which volcanic centres are the most hazardous, where are they, and what is likelihood of activity? Together with a better understanding of which hazard processes pose the most risk at particular distances, a better global understanding of hazard exposure will be possible. An improved global overview of hazardous marine volcanic centres by the volcanology and ocean exploration community is imperative to more adequately inform risk mitigation for shipping industries.

7 CONCLUSIONS

The risks posed to shipping around volcanic centres are varied and generally poorly understood. When considering surface shipping and coastal infrastructure, the risk of a total loss to shipping is relatively low. However, if an eruption does impact shipping, the potential for delays, diversions, and mechanical failures is higher. Given the increasing dependence on global shipping freight and its growth, the time sensitivity of many deliveries, and the costs related to the running of shipping operations, volcanic activity should be a great subject of concern.

Considerable work must be done in establishing baseline values for risk in the offshore setting and in exploring the wider hazards not explored here. However, there is already sufficient knowledge in many areas for policy and available datasets to start addressing some of these hazards in advance of a more substantial crisis.

Given the myriad hazards which may be associated with explosive eruptions, we suggest that a collaborative and meaningful effort is prioritised toward the establishment of a shipping-oriented alert system which runs similarly to, in parallel with, or as part of the Volcanic Ash Advisory Centres designated under the International Civil Aviation Organization. In addition, there are significant steps that can be taken in the volcanology and natural hazards scientific communities to progress research in the areas needed to better inform and validate the proposed policy and regulatory changes.

AUTHOR CONTRIBUTIONS

PC conceived the initial idea for this work together with initial literature review material and conducted a number of experiments while at sea to evidence several arguments made herein (e.g., Figure 5). PR wrote the draft manuscript with regular discussion with SM, which all authors contributed to the editing and revision of. All authors contributed to data collation and figure design and preparation.

ACKNOWLEDGEMENTS

We are very grateful to S. Jenkins for providing the original ashfall recurrence shapefile data which enabled us to produce Figure 4. Useful discussions with Isobel Yeo and Mike Clare have helped inform the direction and content of this review. We are very grateful for the constructive and detailed insights provided by two anonymous reviewers that substantially improved the manuscript.

DATA AVAILABILITY

All data used for generating the GIS outputs is available through open-source channels as referenced in the relevant figure captions. The centroid analysis assessing the number of volcanoes within 100 km of the ports listed in the World Port Index is available here: doi.org/10.5281/zenodo.15365446.

COPYRIGHT NOTICE

© The Author(s) 2026. This article is distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

REFERENCES

- Alexander, D. (2013). "Volcanic ash in the atmosphere and risks for civil aviation: A study in European crisis management". *International Journal of Disaster Risk Science* 4(1), pages 9–19. DOI: [10.1007/s13753-013-0003-0](https://doi.org/10.1007/s13753-013-0003-0).
- Allen, S., R. Fiske, and K. Cashman (2008). "Quenching of steam-charged pumice: Implications for submarine pyroclastic volcanism". *Earth and Planetary Science Letters* 274(1-2), pages 40–49. DOI: [10.1016/j.epsl.2008.06.050](https://doi.org/10.1016/j.epsl.2008.06.050).
- Andronico, D., E. Del Bello, C. D'Oriano, P. Landi, F. Pardini, P. Scarlato, M. de' Michieli Vitturi, J. Taddeucci, A. Cristaldi, F. Ciancitto, F. Pennacchia, T. Ricci, and F. Valentini (2021). "Uncovering the eruptive patterns of the 2019 double paroxysm eruption crisis of Stromboli volcano". *Nature Communications* 12(1). DOI: [10.1038/s41467-021-24420-1](https://doi.org/10.1038/s41467-021-24420-1).
- Aravena, A., A. Tadini, A. Bevilacqua, P. Samaniego, B. Bernard, S. Hidalgo, J.-L. Le Pennec, P. Martínez-Yáñez, J. García, and O. Roche (2024). "Probabilistic, scenario-based hazard assessment for pyroclastic density currents at Tungurahua volcano, Ecuador". *Bulletin of Volcanology* 86(10). DOI: [10.1007/s00445-024-01768-y](https://doi.org/10.1007/s00445-024-01768-y).
- Asano, T. and A. Nagayama (2021). "Analysis of workload required for removal of drifting pumice after a volcanic disaster as an aspect of a port business continuity plan: A case study of Kagoshima Port, Japan". *International Journal of Disaster Risk Reduction* 64, page 102511. DOI: [10.1016/j.ijdrr.2021.102511](https://doi.org/10.1016/j.ijdrr.2021.102511).
- Barsotti, S., D. I. Di Rienzo, T. Thordarson, B. B. Björnsson, and S. Karlsdóttir (2018). "Assessing Impact to Infrastructures Due to Tephra Fallout From Öræfajökull Volcano (Iceland) by Using a Scenario-Based Approach and a Numerical

- Model". *Frontiers in Earth Science* 6. DOI: [10.3389/feart.2018.00196](https://doi.org/10.3389/feart.2018.00196).
- BBC News (2018). *Kilauea volcano: 'Lava bomb' injures 23 on Hawaii boat*. URL: <https://www.bbc.com/news/world-us-canada-44851526>.
- Bebbington, M., S. J. Cronin, I. Chapman, and M. B. Turner (2008). "Quantifying volcanic ash fall hazard to electricity infrastructure". *Journal of Volcanology and Geothermal Research* 177(4), pages 1055–1062. DOI: [10.1016/j.jvolgeores.2008.07.023](https://doi.org/10.1016/j.jvolgeores.2008.07.023).
- Bernard, E. and V. Titov (2015). "Evolution of tsunami warning systems and products". *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 373(2053), page 20140371. DOI: [10.1098/rsta.2014.0371](https://doi.org/10.1098/rsta.2014.0371).
- Bevilacqua, A., A. Aravena, W. Aspinall, A. Costa, S. Mahony, A. Neri, S. Sparks, and B. Hill (2022). "Assessing minimum pyroclastic density current mass to impact critical infrastructures: example from Aso caldera (Japan)". *Natural Hazards and Earth System Sciences* 22(10), pages 3329–3348. DOI: [10.5194/nhess-22-3329-2022](https://doi.org/10.5194/nhess-22-3329-2022).
- Bonaccorso, A., S. Calvari, G. Garfi, L. Lodato, and D. Patanè (2003). "Dynamics of the December 2002 flank failure and tsunami at Stromboli volcano inferred by volcanological and geophysical observations". *Geophysical Research Letters* 30(18). DOI: [10.1029/2003gl017702](https://doi.org/10.1029/2003gl017702).
- Bonadonna, C. and A. Costa (2013). "Plume height, volume, and classification of explosive volcanic eruptions based on the Weibull function". *Bulletin of Volcanology* 75(8). DOI: [10.1007/s00445-013-0742-1](https://doi.org/10.1007/s00445-013-0742-1).
- Borrero, J. C., P. J. Lynett, and N. Kalligeris (2015). "Tsunami currents in ports". *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 373(2053), page 20140372. DOI: [10.1098/rsta.2014.0372](https://doi.org/10.1098/rsta.2014.0372).
- Bougouin, A., R. Paris, and O. Roche (2020). "Impact of Fluidized Granular Flows into Water: Implications for Tsunamis Generated by Pyroclastic Flows". *Journal of Geophysical Research: Solid Earth* 125(5). DOI: [10.1029/2019jb018954](https://doi.org/10.1029/2019jb018954).
- Brown, S. K., R. Sparks, and S. Jenkins (2015). "Global distribution of volcanic threat". *Global Volcanic Hazards and Risk*. Edited by S. C. Loughlin, S. Sparks, S. K. Brown, S. F. Jenkins, and C. Vye-Brown. Cambridge University Press, pages 359–370. ISBN: 9781316276273. DOI: [10.1017/cbo9781316276273.025](https://doi.org/10.1017/cbo9781316276273.025).
- Bryan, S., A. Cook, J. Evans, P. Colls, M. Wells, M. Lawrence, J. Jell, A. Greig, and R. Leslie (2004). "Pumice rafting and faunal dispersion during 2001–2002 in the Southwest Pacific: record of a dacitic submarine explosive eruption from Tonga". *Earth and Planetary Science Letters* 227(1–2), pages 135–154. DOI: [10.1016/j.epsl.2004.08.009](https://doi.org/10.1016/j.epsl.2004.08.009).
- Bryan, S. E., A. G. Cook, J. P. Evans, K. Hebden, L. Hurrey, P. Colls, J. S. Jell, D. Weatherley, and J. Firn (2012). "Rapid, Long-Distance Dispersal by Pumice Rafting". *PLoS ONE* 7(7). Edited by N. Houlie, e40583. DOI: [10.1371/journal.pone.0040583](https://doi.org/10.1371/journal.pone.0040583).
- Bryan, S., A. Clarke, L. Vanderkluyzen, C. Groppi, S. Paine, D. W. Bliss, J. Aberle, and P. Mauskopf (2017). "Measuring Water Vapor and Ash in Volcanic Eruptions With a Millimeter-Wave Radar/Imager". *IEEE Transactions on Geoscience and Remote Sensing* 55(6), pages 3177–3185. DOI: [10.1109/tgrs.2017.2663381](https://doi.org/10.1109/tgrs.2017.2663381).
- Caron, A. (2013). "The Danish West Indies and the Montagne Pelée Eruption of 1902". *Caribbean Genealogy Library*, pages 1–23.
- Cas, R., J. V. Wright, and G. Giordano (2024). "Subaqueous Flows of Pyroclastic Debris and Their Deposits: Origins, and Distinguishing Hot Gas-Supported Flows and Deposits from Water-Supported Ones". *Volcanology*. Edited by R. Cas, G. Giordano, and J. V. Wright. Springer International Publishing, pages 891–981. ISBN: 9783319666136. DOI: [10.1007/978-3-319-66613-6_11](https://doi.org/10.1007/978-3-319-66613-6_11).
- Chang, Y.-L. K., I. M. McIntosh, T. Miyama, and Y. Miyazawa (2023). "Projection of August 2021 pumice dispersion from the Fukutoku-Oka-no-Ba eruption in the western North Pacific". *Scientific Reports* 13(1). DOI: [10.1038/s41598-023-31058-0](https://doi.org/10.1038/s41598-023-31058-0).
- Chrétien, S. and R. Brousse (1989). "Events preceding the great eruption of 8 May, 1902 at Mount Pelée, Martinique". *Journal of Volcanology and Geothermal Research* 38(1–2), pages 67–75. DOI: [10.1016/0377-0273\(89\)90030-9](https://doi.org/10.1016/0377-0273(89)90030-9).
- Clare, M. A., I. A. Yeo, J. Nash, J. E. Hunt, S. Panuve, A. Wilkie, R. Williams, N. Dowey, P. Rowley, J. Barclay, J. Phillips, J. Scarlett, S. Engwell, T. J. Henstock, S. Seabrook, S. Watson, R. Wysoczanski, M. Ribo, S. Cronin, P. J. Talling, M. Cassidy, S. Watt, and R. Robertson (2025). "Volcanic eruptions and the global subsea telecommunications network". *Bulletin of Volcanology* 87(6). DOI: [10.1007/s00445-025-01832-1](https://doi.org/10.1007/s00445-025-01832-1).
- Clare, M. A., I. A. Yeo, S. Watson, R. Wysoczanski, S. Seabrook, K. Mackay, J. E. Hunt, E. Lane, P. J. Talling, E. Pope, S. Cronin, M. Ribó, T. Kula, D. Tappin, S. Henrys, C. de Ronde, M. Urlaub, S. Kutterolf, S. Fonua, S. Panuve, D. Veverka, R. Rapp, V. Kamalov, and M. Williams (2023). "Fast and destructive density currents created by ocean-entering volcanic eruptions". *Science* 381(6662), pages 1085–1092. DOI: [10.1126/science.ad13038](https://doi.org/10.1126/science.ad13038).
- Clarkson, R. J., E. J. Majewicz, and P. Mack (2016). "A re-evaluation of the 2010 quantitative understanding of the effects volcanic ash has on gas turbine engines". *Proceedings of the Institution of Mechanical Engineers, Part G: Journal of Aerospace Engineering* 230(12), pages 2274–2291. DOI: [10.1177/0954410015623372](https://doi.org/10.1177/0954410015623372).
- Davison, C. R. and T. A. Rutke (2014). "Assessment and Characterization of Volcanic Ash Threat to Gas Turbine Engine Performance". *Journal of Engineering for Gas Turbines and Power* 136(8). DOI: [10.1115/1.4026810](https://doi.org/10.1115/1.4026810).
- Denardo, B., L. Pringle, C. DeGrace, and M. McGuire (2001). "When do bubbles cause a floating body to sink?" *American Journal of Physics* 69(10), pages 1064–1072. DOI: [10.1119/1.1383600](https://doi.org/10.1119/1.1383600).
- Di Roberto, A., A. Bertagnini, M. Pompilio, and M. Bisson (2014). "Pyroclastic density currents at Stromboli volcano (Aeolian Islands, Italy): a case study of the 1930 eruption".

- Bulletin of Volcanology* 76(6). DOI: [10.1007/s00445-014-0827-5](https://doi.org/10.1007/s00445-014-0827-5).
- Druitt, T., S. Kutterolf, T. A. Ronge, C. Hübscher, P. Nomikou, J. Preine, R. Gertisser, J. Karstens, J. Keller, O. Koukousioura, M. Manga, A. Metcalfe, M. McCanta, I. McIntosh, K. Pank, A. Woodhouse, S. Beethe, C. Berthod, S. Chiyonobu, H. Chen, A. Clark, S. DeBari, R. Johnston, A. Peccia, Y. Yamamoto, A. Bernard, T. F. Perez, C. Jones, K. B. Joshi, G. Kletetschka, X. Li, A. Morris, P. Polymenakou, M. Tomlinaga, D. Papanikolaou, K.-L. Wang, and H.-Y. Lee (2024). “Giant offshore pumice deposit records a shallow submarine explosive eruption of ancestral Santorini”. *Communications Earth and Environment* 5(1). DOI: [10.1038/s43247-023-01171-z](https://doi.org/10.1038/s43247-023-01171-z).
- Dunn, M. G., A. J. Baran, and J. Miatech (1996). “Operation of Gas Turbine Engines in Volcanic Ash Clouds”. *Journal of Engineering for Gas Turbines and Power* 118(4), pages 724–731. DOI: [10.1115/1.2816987](https://doi.org/10.1115/1.2816987).
- Engwell, S., L. G. Mastin, C. Bonadonna, S. Barsotti, N. I. Deligne, and B. A. Oladottir (2024). “Characterising, quantifying, and accessing eruption source parameters of explosive volcanic eruptions for operational simulation of tephra dispersion: a current view and future perspectives”. *Bulletin of Volcanology* 86(7). DOI: [10.1007/s00445-024-01706-y](https://doi.org/10.1007/s00445-024-01706-y).
- Evans, K. and F. McCoy (2020). “Precursory eruptive activity and implied cultural responses to the Late Bronze Age (LBA) eruption of Thera (Santorini, Greece)”. *Journal of Volcanology and Geothermal Research* 397, page 106868. DOI: [10.1016/j.jvolgeores.2020.106868](https://doi.org/10.1016/j.jvolgeores.2020.106868).
- Fouqué, F. (1879). *Santorin et ses éruption*. G. Masson.
- Freundt, A. (2003). “Entrance of hot pyroclastic flows into the sea: experimental observations”. *Bulletin of Volcanology* 65(2-3), pages 144–164. DOI: [10.1007/s00445-002-0250-1](https://doi.org/10.1007/s00445-002-0250-1).
- Frey, S. E., M. Jutzeler, and R. J. Carey (2026). “A review of the supercritical state of eruption-fed volcanic density currents in subaerial and subaqueous settings”. *Sedimentology*. DOI: [10.1111/sed.70083](https://doi.org/10.1111/sed.70083).
- Furst, S., M. Urlaub, E. Klein, and C. Bonanati (2023). “Are eruptions reliable precursors to marine volcano collapses?” *Frontiers in Earth Science* 11. DOI: [10.3389/feart.2023.1130561](https://doi.org/10.3389/feart.2023.1130561).
- Gass, I. G., P. G. Harris, and M. W. Holdgate (1963). “Pumice Eruption in the Area of the South Sandwich Islands”. *Geological Magazine* 100(4), pages 321–330. DOI: [10.1017/s0016756800056053](https://doi.org/10.1017/s0016756800056053).
- Giachetti, T., R. Paris, K. Kelfoun, and B. Ontowirjo (2012). “Tsunami hazard related to a flank collapse of Anak Krakatau Volcano, Sunda Strait, Indonesia”. *Geological Society, London, Special Publications* 361(1), pages 79–90. DOI: [10.1144/sp361.7](https://doi.org/10.1144/sp361.7).
- Giehl, C., R. A. Brooker, H. Marxer, and M. Nowak (2017). “An experimental simulation of volcanic ash deposition in gas turbines and implications for jet engine safety”. *Chemical Geology* 461, pages 160–170. DOI: [10.1016/j.chemgeo.2016.11.024](https://doi.org/10.1016/j.chemgeo.2016.11.024).
- Giordano, D., A. Nichols, and D. Dingwell (2005). “Glass transition temperatures of natural hydrous melts: a relationship with shear viscosity and implications for the welding process”. *Journal of Volcanology and Geothermal Research* 142(1-2), pages 105–118. DOI: [10.1016/j.jvolgeores.2004.10.015](https://doi.org/10.1016/j.jvolgeores.2004.10.015).
- Giordano, G. and G. De Astis (2020). “The summer 2019 basaltic Vulcanian eruptions (paroxysms) of Stromboli”. *Bulletin of Volcanology* 83(1). DOI: [10.1007/s00445-020-01423-2](https://doi.org/10.1007/s00445-020-01423-2).
- Global Volcanism Program (1994). “Rabaul (Papua New Guinea) Tavurvur remains active; details of September eruptions”. *Bulletin of the Global Volcanism Network* 19:9. Edited by E. Venzke. DOI: [10.5479/si.GVP.BGVN199409-252140](https://doi.org/10.5479/si.GVP.BGVN199409-252140).
- (2025). *Volcanoes of the World, v.5.2.4*. Edited by E. Venzke. DOI: [10.5479/si.GVP.VOTW5-2025.5.3](https://doi.org/10.5479/si.GVP.VOTW5-2025.5.3).
- Gordon, K. D., J. W. Cole, M. D. Rosenberg, and D. M. Johnston (2005). “Effects of Volcanic Ash on Computers and Electronic Equipment”. *Natural Hazards* 34(2), pages 231–262. DOI: [10.1007/s11069-004-1514-1](https://doi.org/10.1007/s11069-004-1514-1).
- Grilli, S. T., D. R. Tappin, S. Carey, S. F. L. Watt, S. N. Ward, A. R. Grilli, S. L. Engwell, C. Zhang, J. T. Kirby, L. Schambach, and M. Muin (2019). “Modelling of the tsunami from the December 22, 2018 lateral collapse of Anak Krakatau volcano in the Sunda Straits, Indonesia”. *Scientific Reports* 9(1). DOI: [10.1038/s41598-019-48327-6](https://doi.org/10.1038/s41598-019-48327-6).
- Guibert, J.-S., M. Guérout, and L. Serra (2018). “Past and Present Research in the Underwater Archaeology of Saint-Pierre, Martinique (FWI)”. *Underwater Archaeology Proceedings*, pages 149–154.
- Hansell, A. and C. Oppenheimer (2004). “Health Hazards from Volcanic Gases: A Systematic Literature Review”. *Archives of Environmental Health: An International Journal* 59(12), pages 628–639. DOI: [10.1080/00039890409602947](https://doi.org/10.1080/00039890409602947).
- Harris, D., W. Rose, R. Roe, and M. Thompson (1981). “Radar observations of ash eruptions”. Lipman, P. and D. Mullineaux. *The 1980 eruptions of Mount St. Helens, Washington*. Professional Paper 1250. U.S. Geological Survey, pages 323–333. DOI: [10.3133/pp1250](https://doi.org/10.3133/pp1250).
- He, S., L. Qi, M. Wang, C. Le, and C. Hu (2025). “Pumice rafts in the global ocean: a remote sensing assessment”. *GIScience and Remote Sensing* 62(1). DOI: [10.1080/15481603.2025.2555093](https://doi.org/10.1080/15481603.2025.2555093).
- Horwell, C. J. and P. J. Baxter (2006). “The respiratory health hazards of volcanic ash: a review for volcanic risk mitigation”. *Bulletin of Volcanology* 69(1), pages 1–24. DOI: [10.1007/s00445-006-0052-y](https://doi.org/10.1007/s00445-006-0052-y).
- Hueschen, M. A. (2010). “Can bubbles sink ships?” *American Journal of Physics* 78(2), pages 139–141. DOI: [10.1119/1.3263819](https://doi.org/10.1119/1.3263819).
- Imamura, F., A. Suppasri, T. Arikawa, S. Koshimura, K. Satake, and Y. Tanioka (2022). “Preliminary Observations and Impact in Japan of the Tsunami Caused by the Tonga Volcanic Eruption on January 15, 2022”. *Pure and Applied Geophysics* 179(5), pages 1549–1560. DOI: [10.1007/s00024-022-03058-0](https://doi.org/10.1007/s00024-022-03058-0).

- International Maritime Organization (2016). *International Code for Ships Operating in Polar Waters (Polar Code)*. URL: <https://www.imo.org/en/ourwork/safety/pages/polar-code.aspx>.
- (2020). *Fourth Greenhouse Gas Study 2020*. URL: <https://wwwcdn.imo.org/localresources/en/OurWork/Environment/Documents/Fourth%20IMO%20GHG%20Study%202020%20-%20Full%20report%20and%20annexes.pdf>.
- Iskandar, M. R., Y.-G. Park, K. Kim, H. Jin, S. Seo, and Y. H. Kim (2023). “Tracking the pumice rafts from the Fukutoku-Okanoba submarine volcano with Satellites and a Lagrangian Particles trajectory model”. *Marine Pollution Bulletin* 193, page 115254. DOI: [10.1016/j.marpolbul.2023.115254](https://doi.org/10.1016/j.marpolbul.2023.115254).
- Jaggard, T. (1949). “Fourth special report of the Hawaiian Volcano Observatory of the U.S. Geological Survey and the Hawaiian Volcano Research Association: Steam blast volcanic eruptions: A study of Mount Pelée in Martinique as type volcano”. Edited by R. Finch. Report of the Hawaiian Volcano Observatory. U.S. Geological Survey, pages 1–137.
- Jenkins, S., T. Wilson, C. Magill, V. Miller, C. Stewart, R. Blong, W. Marzocchi, M. Boulton, C. Bonadonna, and A. Costa (2015). “Volcanic ash fall hazard and risk”. *Global Volcanic Hazards and Risk*. Edited by S. Loughlin, S. Sparks, S. Brown, S. Jenkin, and C. Vye-Brown. Cambridge University Press, pages 173–222. ISBN: 9781316276273. DOI: [10.1017/cbo9781316276273.005](https://doi.org/10.1017/cbo9781316276273.005).
- Jenkins, S. F., A. McSporran, T. M. Wilson, C. Stewart, G. Leonard, S. Cevuard, and E. Garaebiti (2024). “Tephra fall impacts to buildings: the 2017–2018 Manaro Voui eruption, Vanuatu”. *Frontiers in Earth Science* 12. DOI: [10.3389/feart.2024.1392098](https://doi.org/10.3389/feart.2024.1392098).
- Johnson, R. and N. Threlfall (1985). *Volcano Town: The 1937-43 Eruptions at Rabaul*. Robert Brown & Associates. ISBN: 978-1760466039.
- Jutzeler, M., M. Manga, J. White, P. Talling, A. Proussevitch, S. Watt, M. Cassidy, R. Taylor, A. Le Friant, and O. Ishizuka (2017). “Submarine deposits from pumiceous pyroclastic density currents traveling over water: An outstanding example from offshore Montserrat (IODP 340)”. *Geological Society of America Bulletin* 129(3-4), pages 392–414. DOI: [10.1130/b31448.1](https://doi.org/10.1130/b31448.1).
- Jutzeler, M., R. Marsh, R. J. Carey, J. D. L. White, P. J. Talling, and L. Karlstrom (2014). “On the fate of pumice rafts formed during the 2012 Havre submarine eruption”. *Nature Communications* 5(1). DOI: [10.1038/ncomms4660](https://doi.org/10.1038/ncomms4660).
- Jutzeler, M., R. Marsh, E. van Sebille, T. Mittal, R. J. Carey, K. E. Fauria, M. Manga, and J. McPhie (2020). “Ongoing Dispersal of the 7 August 2019 Pumice Raft From the Tonga Arc in the Southwestern Pacific Ocean”. *Geophysical Research Letters* 47(5). DOI: [10.1029/2019gl086768](https://doi.org/10.1029/2019gl086768).
- Karátson, D., T. Telbisz, R. Gertisser, T. Strasser, P. Nomikou, T. Druitt, V. Vereb, X. Quidelleur, and S. Kósik (2020). “Constraining the landscape of Late Bronze Age Santorini prior to the Minoan eruption: Insights from volcanological, geomorphological and archaeological findings”. *Journal of Volcanology and Geothermal Research* 401, page 106911. DOI: [10.1016/j.jvolgeores.2020.106911](https://doi.org/10.1016/j.jvolgeores.2020.106911).
- Kubo, M., I.-s. Cho, S. Sakakibara, E. Kobayashi, and S. Koshimura (2005). “The Influence of Tsunamis on Moored Ships and Ports”. *Journal of Navigation and Port Research* 29, pages 319–325.
- Lacasse, C., S. Karlsdóttir, G. Larsen, H. Soosalu, W. I. Rose, and G. G. J. Ernst (2004). “Weather radar observations of the Hekla 2000 eruption cloud, Iceland”. *Bulletin of Volcanology* 66(5), pages 457–473. DOI: [10.1007/s00445-003-0329-3](https://doi.org/10.1007/s00445-003-0329-3).
- Le Friant, A., C. Deplus, G. Boudon, R. Sparks, J. Trofimovs, and P. Talling (2009). “Submarine deposition of volcanoclastic material from the 1995–2005 eruptions of Soufrière Hills volcano, Montserrat”. *Journal of the Geological Society* 166(1), pages 171–182. DOI: [10.1144/0016-76492008-047](https://doi.org/10.1144/0016-76492008-047).
- Lee, J. M.-y. and E. Y.-c. Wong (2021). “Suez Canal blockage: an analysis of legal impact, risks and liabilities to the global supply chain”. *MATEC Web of Conferences* 339. Edited by P. Khvedelidze, B. Gechbaia, and K. Goletiani, page 01019. DOI: [10.1051/mateconf/202133901019](https://doi.org/10.1051/mateconf/202133901019).
- Legros, F. and T. Druitt (2000). “On the emplacement of ignimbrite in shallow-marine environments”. *Journal of Volcanology and Geothermal Research* 95(1-4), pages 9–22. DOI: [10.1016/s0377-0273\(99\)00116-x](https://doi.org/10.1016/s0377-0273(99)00116-x).
- Li, R. (2023). “The Impact of Transportation Delays on the Supply Chain — Using the Suez Canal as an Example”. *Proceedings of the 2022 2nd International Conference on Business Administration and Data Science (BADs 2022)*. Edited by V. Fernández-Viagas, B. K. Kandel, C. Valmohammadi, and H. Mallick. Atlantis Press International BV, pages 899–905. ISBN: 9789464631029. DOI: [10.2991/978-94-6463-102-9_93](https://doi.org/10.2991/978-94-6463-102-9_93).
- Lipiejko, N., C. N. Whittaker, E. M. Lane, J. D. L. White, and W. L. Power (2022). “Experimental Modeling of Tsunamis Generated by Pyroclastic Density Currents: The Effects of Particle Size Distribution on Wave Generation”. *Journal of Geophysical Research: Solid Earth* 127(11). DOI: [10.1029/2022jb024847](https://doi.org/10.1029/2022jb024847).
- Liupeng, J., W. Guangsheng, F. Xuejun, Y. Tong, and L. Zhiyi (2024). “Study on cascading failure vulnerability of the 21st-century Maritime Silk Road container shipping network”. *Journal of Transport Geography* 117, page 103891. DOI: [10.1016/j.jtrangeo.2024.103891](https://doi.org/10.1016/j.jtrangeo.2024.103891).
- López-Saavedra, M. and J. Martí (2023). “Reviewing the multi-hazard concept. Application to volcanic islands”. *Earth-Science Reviews* 236, page 104286. DOI: [10.1016/j.earscirev.2022.104286](https://doi.org/10.1016/j.earscirev.2022.104286).
- Loughlin, S., R. Sparks, S. Brown, S. Jenkins, and C. Vye-Brown (2015). *Global Volcanic Hazards and Risk*. Cambridge University Press. ISBN: 9781316276273. DOI: [10.1017/cbo9781316276273](https://doi.org/10.1017/cbo9781316276273).
- Lynett, P., M. McCann, Z. Zhou, W. Renteria, J. Borrero, D. Greer, O. Fa’anunu, C. Bosserelle, B. Jaffe, S. La Selle, A. Ritchie, A. Snyder, B. Nasr, J. Bott, N. Graehl, C. Synolakis, B. Ebrahimi, and G. E. Cinar (2022). “Diverse tsunamigenesis triggered by the Hunga Tonga-Hunga Ha’apai eruption”.

- Nature* 609(7928), pages 728–733. DOI: [10.1038/s41586-022-05170-6](https://doi.org/10.1038/s41586-022-05170-6).
- Maeno, F. (2025). “Global database on surface phenomena and hazards of explosive submarine eruptions with application to hazards of the Kolumbo Volcanic Field”. *Bulletin of Volcanology* 87(10). DOI: [10.1007/s00445-025-01871-8](https://doi.org/10.1007/s00445-025-01871-8).
- Mani, L., A. Tzachor, and P. Cole (2021). “Global catastrophic risk from lower magnitude volcanic eruptions”. *Nature Communications* 12(1). DOI: [10.1038/s41467-021-25021-8](https://doi.org/10.1038/s41467-021-25021-8).
- Martin, T. R., A. P. Wehner, and J. Butler (1986). “Chapter 7: Evaluation of Physical Health Effects Due to Volcanic Hazards: The Use of Experimental Systems to Estimate the Pulmonary Toxicity of Volcanic Ash”. *American Journal of Public Health* 76(Suppl), pages 59–65. DOI: [10.2105/ajph.76.suppl.59](https://doi.org/10.2105/ajph.76.suppl.59).
- Marzano, F. S., E. Picciotti, M. Montopoli, and G. Vulpiani (2013). “Inside Volcanic Clouds: Remote Sensing of Ash Plumes Using Microwave Weather Radars”. *Bulletin of the American Meteorological Society* 94(10), pages 1567–1586. DOI: [10.1175/bams-d-11-00160.1](https://doi.org/10.1175/bams-d-11-00160.1).
- Marzocchi, W. and G. Woo (2009). “Principles of volcanic risk metrics: Theory and the case study of Mount Vesuvius and Campi Flegrei, Italy”. *Journal of Geophysical Research: Solid Earth* 114(B3). DOI: [10.1029/2008jb005908](https://doi.org/10.1029/2008jb005908).
- McGuire, W. (2003). “Volcano instability and lateral collapse”. *Revista* 1, pages 33–45.
- Milia, A., M. M. Torrente, and A. Zuppetta (2003). “Offshore debris avalanches at Somma–Vesuvius volcano (Italy): implications for hazard evaluation”. *Journal of the Geological Society* 160(2), pages 309–317. DOI: [10.1144/0016-764902-045](https://doi.org/10.1144/0016-764902-045).
- Minakami, T. (1956). “Report on volcanic activities and volcanological studies in Japan for the period from 1951 to 1954”. *Bulletin Volcanologique* 18(1), pages 39–76. DOI: [10.1007/bf02596612](https://doi.org/10.1007/bf02596612).
- Mitchell, S. J., K. E. Fauria, B. F. Houghton, and R. J. Carey (2021). “Sink or float: microtextural controls on the fate of pumice deposition during the 2012 submarine Havre eruption”. *Bulletin of Volcanology* 83(11). DOI: [10.1007/s00445-021-01497-6](https://doi.org/10.1007/s00445-021-01497-6).
- Morimoto, R. and J. Ohsaka (1970). “The 1970-activity of Myojin reef”. *Journal of Geography (Chigaku Zasshi)* 79(6), pages 301–320. DOI: [10.5026/jgeography.79.6_301](https://doi.org/10.5026/jgeography.79.6_301).
- Muhari, A., I. Charvet, F. Tsuyoshi, A. Suppasri, and F. Ima-mura (2015). “Assessment of tsunami hazards in ports and their impact on marine vessels derived from tsunami models and the observed damage data”. *Natural Hazards* 78(2), pages 1309–1328. DOI: [10.1007/s11069-015-1772-0](https://doi.org/10.1007/s11069-015-1772-0).
- Musée Frank A Perret (2019-). *Mémorial de la Catastrophe de 1902 [Exhibition]*. St. Pierre, Martinique.
- Nagai, M. and S. Nakada (2022). “Impact on the Electric Infrastructure Due to Volcanic Ash from a Hydrovolcanic Eruption of Aso Volcano in 2016”. *Journal of Disaster Research* 17(5), pages 829–838. DOI: [10.20965/jdr.2022.p0829](https://doi.org/10.20965/jdr.2022.p0829).
- Nagao, R., T. Yokoyama, and S. Takeuchi (2025). “Pumice flotation time: how does the infiltration of water progress inside pumice?” *Progress in Earth and Planetary Science* 12(1). DOI: [10.1186/s40645-025-00750-2](https://doi.org/10.1186/s40645-025-00750-2).
- National Geospatial-Intelligence Agency (2022). *World Port Index (WPI)*. URL: <https://www.reuters.com/business/energy/oil-prices-rise-further-monday-mideast-war-escalates-2026-03-22/>.
- Nishikawa, H., T. Kuwatani, N. Tada, and H. Kayama Watanabe (2023). “Simulated distributions of pumice rafts in Japan following eruptions at volcanic islands and submarine volcanoes”. *Progress in Earth and Planetary Science* 10(1). DOI: [10.1186/s40645-023-00552-4](https://doi.org/10.1186/s40645-023-00552-4).
- Nomikou, P., S. Carey, D. Papanikolaou, K. Croff Bell, D. Sakellariou, M. Alexandri, and K. Bejelou (2012). “Submarine volcanoes of the Kolumbo volcanic zone NE of Santorini Caldera, Greece”. *Global and Planetary Change* 90-91, pages 135–151. DOI: [10.1016/j.gloplacha.2012.01.001](https://doi.org/10.1016/j.gloplacha.2012.01.001).
- Oze, C., J. Cole, A. Scott, T. Wilson, G. Wilson, S. Gaw, S. Hampton, C. Doyle, and Z. Li (2014). “Corrosion of metal roof materials related to volcanic ash interactions”. *Natural Hazards* 71(1), pages 785–802. DOI: [10.1007/s11069-013-0943-0](https://doi.org/10.1007/s11069-013-0943-0).
- Paris, R. (2015). “Source mechanisms of volcanic tsunamis”. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 373(2053), page 20140380. DOI: [10.1098/rsta.2014.0380](https://doi.org/10.1098/rsta.2014.0380).
- Prata, A. J. and A. Tupper (2009). “Aviation hazards from volcanoes: the state of the science”. *Natural Hazards* 51(2), pages 239–244. DOI: [10.1007/s11069-009-9415-y](https://doi.org/10.1007/s11069-009-9415-y).
- Pratson, L. F. (2023). “Assessing impacts to maritime shipping from marine chokepoint closures”. *Communications in Transportation Research* 3, page 100083. DOI: [10.1016/j.commtr.2022.100083](https://doi.org/10.1016/j.commtr.2022.100083).
- Pyle, D. M. (1989). “The thickness, volume and grainsize of tephra fall deposits”. *Bulletin of Volcanology* 51(1), pages 1–15. DOI: [10.1007/bf01086757](https://doi.org/10.1007/bf01086757).
- Ramadhani, R. and A. Marzaman (2024). “Maritime stability in the Strait of Hormuz: Challenges, global impacts, and multilateral diplomacy”. *HYPOTHESIS: Multidisciplinary Journal Of Social Sciences* 3(02), pages 81–96. DOI: [10.62668/hypothesis.v3i02.1293](https://doi.org/10.62668/hypothesis.v3i02.1293).
- Redick, N. R. (2023). “Review of Pumice Raft Formation Environments, Saturation, and Dispersal Mechanisms”. *McGill Science Undergraduate Research Journal* 18(1), B19–B25. DOI: [10.26443/msurj.v18i1.187](https://doi.org/10.26443/msurj.v18i1.187).
- Resing, J. A. and F. J. Sansone (1999). “The chemistry of lava–seawater interactions: the generation of acidity”. *Geochimica et Cosmochimica Acta* 63(15), pages 2183–2198. DOI: [10.1016/s0016-7037\(99\)00193-3](https://doi.org/10.1016/s0016-7037(99)00193-3).
- Reuters (2026). *Oil prices to rise further on Monday as Mideast war escalates*. URL: <https://www.reuters.com/business/energy/oil-prices-rise-further-monday-mideast-war-escalates-2026-03-22/>.
- Ripepe, M. and G. Lacanna (2024). “Volcano generated tsunami recorded in the near source”. *Nature Communications* 15(1). DOI: [10.1038/s41467-024-45937-1](https://doi.org/10.1038/s41467-024-45937-1).
- Salgado, P., G. Villarosa, D. Beigt, V. Outes, C. Stewart, and F. Baranzini (2023). “Impacts on water transport networks after three widespread volcanic ashfalls in Andean Patago-

- nian lakes". *Volcanica* 6(2), pages 173–200. DOI: [10.30909/vol.06.02.173200](https://doi.org/10.30909/vol.06.02.173200).
- Salgado, P. A., G. Villarosa, D. Beigt, and V. Outes (2022). "Water evacuations in remote tourist regions: evaluating case studies from natural hazards in North Patagonian lakes, Argentina". *Journal of Mountain Science* 19(6), pages 1782–1807. DOI: [10.1007/s11629-021-7207-3](https://doi.org/10.1007/s11629-021-7207-3).
- Sardain, A., E. Sardain, and B. Leung (2019). "Global forecasts of shipping traffic and biological invasions to 2050". *Nature Sustainability* 2(4), pages 274–282. DOI: [10.1038/s41893-019-0245-y](https://doi.org/10.1038/s41893-019-0245-y).
- Sarna-Wojcicki, A. M., S. Shipley, R. Waitt, D. Dzurisin, W. Hays, J. Davis, S. Wood, and T. Baderidge (1980). "Areal distribution, thickness, and volume of downwind ash from the May 18, 1980, eruption of Mount St. Helens". Open-File Report 80-1078. U.S. Geological Survey. DOI: [10.3133/ofr801078](https://doi.org/10.3133/ofr801078).
- Satow, C., S. Watt, M. Cassidy, D. Pyle, and Y. N. Deng (2023). "The Contributions of Marine Sediment Cores to Volcanic Hazard Assessments: Present Examples and Future Perspectives". *Geosciences* 13(4), page 124. DOI: [10.3390/geosciences13040124](https://doi.org/10.3390/geosciences13040124).
- Scarth, A. (2002). *La Catastrophe: The Eruption of Mount Pelee, the Worst Volcanic Disaster of the 20th Century*. Oxford University Press. ISBN: 978-0195218398.
- Seabrook, S., K. Mackay, S. J. Watson, M. A. Clare, J. E. Hunt, I. A. Yeo, E. M. Lane, M. R. Clark, R. Wysoczanski, A. A. Rowden, T. Kula, L. J. Hoffmann, E. Armstrong, and M. J. M. Williams (2023). "Volcaniclastic density currents explain widespread and diverse seafloor impacts of the 2022 Hunga Volcano eruption". *Nature Communications* 14(1). DOI: [10.1038/s41467-023-43607-2](https://doi.org/10.1038/s41467-023-43607-2).
- Shevchenko, A. V., V. N. Dvigalo, T. R. Walter, R. Mania, F. Maccaferri, I. Y. Svirid, A. B. Belousov, and M. G. Belousova (2020). "The rebirth and evolution of Bezymianny volcano, Kamchatka after the 1956 sector collapse". *Communications Earth and Environment* 1(1). DOI: [10.1038/s43247-020-00014-5](https://doi.org/10.1038/s43247-020-00014-5).
- Shifler, D. A. and S. R. Choi (2018). "CMAS Effects on Ship Gas-Turbine Components/Materials". *Volume 1: Aircraft Engine; Fans and Blowers; Marine*. GT2018. American Society of Mechanical Engineers. DOI: [10.1115/gt2018-75865](https://doi.org/10.1115/gt2018-75865).
- Simkin, T. and R. S. Fiske (1983). "KRAKATAU 1883: A Centennial Retrospective on the Eruption and its Atmospheric Effects". *Weatherwise* 36(5), pages 244–254. DOI: [10.1080/00431672.1983.9930158](https://doi.org/10.1080/00431672.1983.9930158).
- Song, W., Y. Lavallée, K.-U. Hess, U. Kueppers, C. Cimarelli, and D. B. Dingwell (2016). "Volcanic ash melting under conditions relevant to ash turbine interactions". *Nature Communications* 7(1). DOI: [10.1038/ncomms10795](https://doi.org/10.1038/ncomms10795).
- Spence, R. J. S., I. Kelman, P. J. Baxter, G. Zuccaro, and S. Petrazzuoli (2005). "Residential building and occupant vulnerability to tephra fall". *Natural Hazards and Earth System Sciences* 5(4), pages 477–494. DOI: [10.5194/nhess-5-477-2005](https://doi.org/10.5194/nhess-5-477-2005).
- Stewart, C., D. E. Damby, C. J. Horwell, T. Elias, E. Iljin-skaya, I. Tomašek, B. M. Longo, A. Schmidt, H. K. Carlsen, E. Mason, P. J. Baxter, S. Cronin, and C. Witham (2022). "Volcanic air pollution and human health: recent advances and future directions". *Bulletin of Volcanology* 84(1). DOI: [10.1007/s00445-021-01513-9](https://doi.org/10.1007/s00445-021-01513-9).
- Sword-Daniels, V., T. M. Wilson, S. Sargeant, T. Rossetto, J. Twigg, D. M. Johnston, S. C. Loughlin, and P. D. Cole (2014). "Chapter 26 Consequences of long-term volcanic activity for essential services in Montserrat: challenges, adaptations and resilience". *Geological Society, London, Memoirs* 39(1), pages 471–488. DOI: [10.1144/m39.26](https://doi.org/10.1144/m39.26).
- Takeuchi, S., K. Ishige, S. Uesawa, and Y. Suwa (2024). "Unsinkable, long-drifting, millimeter-sized pumice of the 2021 eruption of Fukutoku-Oka-no-Ba submarine volcano". *Progress in Earth and Planetary Science* 11(1). DOI: [10.1186/s40645-024-00652-9](https://doi.org/10.1186/s40645-024-00652-9).
- Tanguy, J.-C. (1994). "The 1902–1905 eruptions of Montagne Pelée, Martinique: anatomy and retrospection". *Journal of Volcanology and Geothermal Research* 60(2), pages 87–107. DOI: [10.1016/0377-0273\(94\)90064-7](https://doi.org/10.1016/0377-0273(94)90064-7).
- Terry, J. P., J. Goff, N. Winspear, V. P. Bongolan, and S. Fisher (2022). "Tonga volcanic eruption and tsunami, January 2022: globally the most significant opportunity to observe an explosive and tsunamigenic submarine eruption since AD 1883 Krakatau". *Geoscience Letters* 9(1). DOI: [10.1186/s40562-022-00232-z](https://doi.org/10.1186/s40562-022-00232-z).
- The Guardian (2022). *Tsunami from Tonga volcano eruption leaves trail of flood damage*. URL: <https://www.theguardian.com/world/2022/jan/16/tonga-volcano-eruption-and-tsunami-leaves-trail-of-flood-damage>.
- Thorarinsson, S. (1950). "The eruption of Mt. Hekla 1947 – 1948". *Bulletin Volcanologique* 10(1), pages 157–168. DOI: [10.1007/bf02596085](https://doi.org/10.1007/bf02596085).
- Thouret, J.-C., N. Aisyah, S. F. Jenkins, E. de Bélizal, Sulistiyani, S. J. Charbonnier, D. S. Sayudi, I. G. M. A. Nandaka, G. Mainsant, and A. Solikhin (2023). "Merapi's Lahars: Characteristics, Behaviour, Monitoring, Impact, Hazard Modelling and Risk Assessment". *Merapi Volcano: Geology, Eruptive Activity, and Monitoring of a High-Risk Volcano*. Edited by R. Gertisser, V. Troll, T. Walter, I. Nandaka, and A. Ratdomopurbo. Springer International Publishing, pages 501–552. ISBN: 9783031150401. DOI: [10.1007/978-3-031-15040-1_17](https://doi.org/10.1007/978-3-031-15040-1_17).
- Trofimovs, J., C. Foster, R. S. J. Sparks, S. Loughlin, A. Le Friant, C. Deplus, L. Porritt, T. Christopher, R. Luckett, P. J. Talling, M. R. Palmer, and T. Le Bas (2012). "Submarine pyroclastic deposits formed during the 20th May 2006 dome collapse of the Soufrière Hills Volcano, Montserrat". *Bulletin of Volcanology* 74(2), pages 391–405. DOI: [10.1007/s00445-011-0533-5](https://doi.org/10.1007/s00445-011-0533-5).
- UK Hydrographic Office (2009). *ADMIRALTY Mariner's Handbook (NP100)*.
- UK National Archives (1998). *The Merchant Shipping (Load Line) Regulations 1998, No 2241*. URL: <https://www.legislation.gov.uk/uksi/1998/2241/made>.
- United Nations Trade and Development (2023). *Review of Maritime Transport 2023*. URL: https://unctad.org/system/files/official-document/rmt2023_en.pdf.

- University of the West Indies Seismic Research Centre (n.d.[a]). *Kick 'Em Jenny Yachting Information*. URL: <https://uwiseismic.com/volcanoes/kick-em-jenny/kick-em-jenny-yachting-information/>.
- (n.d.[b]). *Kick-'em-Jenny Hazards*. URL: <https://uwiseismic.com/volcanoes/kick-em-jenny/kej-hazards/>.
- Verolino, A., S. F. Wee, S. F. Jenkins, F. Costa, and A. D. Switzer (2024). “SEATANI: hazards from seamounts in Southeast Asia, Taiwan, and Andaman and Nicobar Islands (eastern India)”. *Natural Hazards and Earth System Sciences* 24(4), pages 1203–1222. DOI: [10.5194/nhess-24-1203-2024](https://doi.org/10.5194/nhess-24-1203-2024).
- Walding, N., R. Williams, N. Dowey, P. Rowley, M. Thomas, S. Osman, M. Johnson, and D. R. Parsons (2025). “The influence of moisture on ash strength: implications for understanding volcanic stratigraphy”. *Bulletin of Volcanology* 87(6). DOI: [10.1007/s00445-025-01821-4](https://doi.org/10.1007/s00445-025-01821-4).
- Walding, N., R. Williams, A. J. Jamieson, and H. A. Stewart (2026). “Deep seafloor exposures of subaerial volcanic lithofacies on the Tonga trench slope: Ground-truthing forearc structure”. *Journal of Volcanology and Geothermal Research* 473, page 108601. DOI: [10.1016/j.jvolgeores.2026.108601](https://doi.org/10.1016/j.jvolgeores.2026.108601).
- Walding, N., R. Williams, P. Rowley, and N. Dowey (2023). “Cohesional behaviours in pyroclastic material and the implications for deposit architecture”. *Bulletin of Volcanology* 85(11). DOI: [10.1007/s00445-023-01682-9](https://doi.org/10.1007/s00445-023-01682-9).
- Walsh, C., N.-J. Lazarou, M. Traut, J. Price, C. Raucci, M. Sharmina, P. Agnolucci, S. Mander, P. Gilbert, K. Anderson, A. Larkin, and T. Smith (2019). “Trade and trade-offs: Shipping in changing climates”. *Marine Policy* 106, page 103537. DOI: [10.1016/j.marpol.2019.103537](https://doi.org/10.1016/j.marpol.2019.103537).
- Ward, S. N. and S. Day (2003). “Ritter Island Volcano-lateral collapse and the tsunami of 1888”. *Geophysical Journal International* 154(3), pages 891–902. DOI: [10.1046/j.1365-246x.2003.02016.x](https://doi.org/10.1046/j.1365-246x.2003.02016.x).
- Wardman, J. B., T. M. Wilson, P. S. Bodger, J. W. Cole, and C. Stewart (2012). “Potential impacts from tephra fall to electric power systems: a review and mitigation strategies”. *Bulletin of Volcanology* 74(10), pages 2221–2241. DOI: [10.1007/s00445-012-0664-3](https://doi.org/10.1007/s00445-012-0664-3).
- Waythomas, C. F. and R. B. Waitt (1998). “Preliminary volcano-hazard assessment for Augustine Volcano, Alaska”. Open-File Report 98-106. U.S. Geological Survey. DOI: [10.3133/ofr98106](https://doi.org/10.3133/ofr98106).
- Whiteside, A., C. Dupouy, A. Singh, R. Frouin, C. Menkes, and J. Lefèvre (2021). “Automatic Detection of Optical Signatures within and around Floating Tonga-Fiji Pumice Rafts Using MODIS, VIIRS, and OLCI Satellite Sensors”. *Remote Sensing* 13(3), page 501. DOI: [10.3390/rs13030501](https://doi.org/10.3390/rs13030501).
- Williams, G. T., S. F. Jenkins, D. W. J. Lee, and S. J. Wee (2021). “How rainfall influences tephra fall loading — an experimental approach”. *Bulletin of Volcanology* 83(6). DOI: [10.1007/s00445-021-01465-0](https://doi.org/10.1007/s00445-021-01465-0).
- Williams, R., P. Rowley, and M. C. Garthwaite (2019). “Reconstructing the Anak Krakatau flank collapse that caused the December 2018 Indonesian tsunami”. *Geology* 47(10), pages 973–976. DOI: [10.1130/g46517.1](https://doi.org/10.1130/g46517.1).
- Wilson, G., T. Wilson, J. Cole, and C. Oze (2012a). “Vulnerability of laptop computers to volcanic ash and gas”. *Natural Hazards* 63(2), pages 711–736. DOI: [10.1007/s11069-012-0176-7](https://doi.org/10.1007/s11069-012-0176-7).
- Wilson, T. M., C. Stewart, V. Sword-Daniels, G. S. Leonard, D. M. Johnston, J. W. Cole, J. Wardman, G. Wilson, and S. T. Barnard (2012b). “Volcanic ash impacts on critical infrastructure”. *Physics and Chemistry of the Earth, Parts A/B/C* 45-46, pages 5–23. DOI: [10.1016/j.pce.2011.06.006](https://doi.org/10.1016/j.pce.2011.06.006).
- World Bank (2022). *The January 15, 2022 Hunga Tonga-Hunga Ha’apai eruption and tsunami, Tonga [Global Rapid Post Disaster Damage Estimation (GRADE) Report]*. International Bank for Reconstruction and Development/The World Bank. URL: <https://thedocs.worldbank.org/en/doc/b69af83e486aa652d4232276ad698c7b-0070062022/original/GRADE-Report-Tonga-Volcanic-Eruption.pdf>.
- World Bank Group (2021). *Data Catalog: Global Shipping Traffic Density*. URL: <https://datacatalog.worldbank.org/search/dataset/0037580/global-shipping-traffic-density>.
- (2023). *Data Catalog: Global - International Ports*. URL: <https://datacatalog.worldbank.org/search/dataset/0038118/global-international-ports>.
- Xu, K., Y. Cui, and J. Liu (2024). “Modeling the resilience of liner shipping network under cascading effects: Considering distance constraints and transportation time”. *Computers and Industrial Engineering* 197, page 110559. DOI: [10.1016/j.cie.2024.110559](https://doi.org/10.1016/j.cie.2024.110559).
- Xu, X., Y. Zhu, M. Xu, W. Deng, and Y. Zuo (2022). “Vulnerability analysis of the global liner shipping network: from static structure to cascading failure dynamics”. *Ocean and Coastal Management* 229, page 106325. DOI: [10.1016/j.ocecoaman.2022.106325](https://doi.org/10.1016/j.ocecoaman.2022.106325).
- Yoshida, K., Y. Tamura, T. Sato, T. Hanyu, Y. Usui, Q. Chang, and S. Ono (2022). “Variety of the drift pumice clasts from the 2021 Fukutoku-Oka-no-Ba eruption, Japan”. *Island Arc* 31(1). DOI: [10.1111/iar.12441](https://doi.org/10.1111/iar.12441).
- Zebrowski, E. (2002). *The Last Days of St. Pierre: The Volcanic Disaster That Claimed 30,000 Live*. Rutgers University Press. ISBN: 978-0813530413.
- Zheng, M., T. Mittal, K. E. Fauria, A. Subramaniam, and M. Jutzeler (2022). “Pumice Raft Detection Using Machine-Learning on Multispectral Satellite Imagery”. *Frontiers in Earth Science* 10. DOI: [10.3389/feart.2022.838532](https://doi.org/10.3389/feart.2022.838532).